INDEX

Acaricidal properties of hexachlorophenol, 839
Acaricides
  concentrated, 363
  orchard, comparative effectiveness of, 857
  use of sleeves for evaluating clothing treatments, 878
Acrobasis caryae, 357
Acrylonitrile-carbon tetrachloride mixture, effect on seeds, 646
Aëdes
  aegypti, 891, 893
  mosquitoes and DDT, 835
Aerosol particles, deposition of, 591
Agonoderus comma, control of, 801
Alaska
  biting insect problems in, 451
  black fly larvae control in, 100
Alaskan mosquito control, 106
Aleurocanthus woglumi, 708
Alfalfa caterpillar
  chemical control of, in California, 487
  polyhedrosis virus against, 301
Alfalfa
  insect pests in N. C., 694
  insecticides against spittlebug on, 653
  insects, control by organic insecticides, 496
  Lygus spp. on, host plant sources of, 93
  snout beetle control, 311
  weevil, new insecticides for, 554
Aliphatic N-substituted p-Nitrobenzamides against codling moth, 151
Allodermanyssus sanguineus, new locality records for, 414
Amblyomma americanum, 716, 880
Anarsia lineatella, 22
Anomala orientalis, 366
Anopheles
  albimanus, fourth instar larva of, 834
  mosquitoes, control of by DDT, 447
Anthonomus
  grandis, 551, 682, 685
  signatus, 559
Aphid
  collection at Tallulah, La., 993
  control and DDT dusts on potato, 766
Aphids
  flue-cured tobacco, 59
  technique for mounting, 846
  tobacco, control of, 544
  turnip, control, 73
Aphis gossypii, 551
Apricots, green fruitworm on, 754
Archips fumiferana, 221
Argyrotaenia citrana, 37, 543, 911
Argyrotaenia velutinana, 354, 398, 629, 701
Asparagus, thrips on, 849
Atta mexicana, 545

Baits
  codling moth, processing aromatic chemicals as, 850
  Japanese beetle, 146
Bamboo, powder-post beetle in, 963
Barber, George W., obituary, 163
Bassus diversus, parasite of Oriental fruit moth, 540
Beans, seed-corn maggot on, 77
Beech
  bark disease and beech scale, 226
  scale and beech bark disease, 226
Beehive, humidity and temperature within, 528
Beet leafhoppers, effect of curly-top-infective, on cantaloup plants, 897
Benzene hexachloride
  absorption of, in poultry, 980
  for plum curculio, 514
  -flavored peaches, 774
  for southern corn rootworm, 992
Bermuda grass seed insects in Arizona, 555
Biological assay method, 987
Bis methanes, toxicity of, to two-spotted spider mite, etc., 993
Black fly
  larvae control
  in Alaska, 100
  by DDT, 392
Black widow spider, habits and control of, 700
Body louse, evaluation of materials against, 694
Boll weevil control, 682, 685, 795
Boxelder bug, control of, 994
Brachyrhinus ligustici, 311
Brazil, livestock parasite control in, 276
Brevicoryne brassicae, 156
Broad-horned flour beetle, 229
Bruchus pisorum, 746
Bud moth control with parathion, 153
Butoxypolypropylene glycol compounds as fly repellents, 281

Cabbage aphid control, 156
Calf starter pellets infested by drug store beetle, 371
California, control of alfalfa caterpillar in, 487
Callitroga americana, 684
Cantaloupe mosaic and insect vectors of, 770
Cantaloupe plants, effects of curly-top-infective beet leafhoppers on, 897
Carpocapsa pomonella, 736
Cat-facing, control of, on peaches, 335
Cattle grub
  control, 837
    by rotenone formulations, 884
  distribution in California, 842
Chalcodermus aeneus, 856
Cherry fruit fly control, 702
Chesnut weevils, DDT for, 825
Chlordane for
  plum curculio, 514
  potato flea beetles, 470
  termites, 541
Chlorinated
  compounds for cowpea insects, 991
  hydrocarbons
    for boll weevil, 685
    for grasshopper control, 135
    against horn fly, 143
    for sheep tick, 410
  hydrocarbon residues, temperatures, and effect on house flies, 891
  organic compounds, insecticidal properties of, 127
Chorioptes bovis, 444
Chrysopa californica, colonization of, against mealybugs, 560
Chrysops vittata, 269
Circulifer tenellus, 897
Cirphis unipuncta, 612
Citrus
  blackfly, control of, 708
  long-tailed mealybug on, 777
  trees, residues on, 829

Clover root borer control, 315
Cnephasia longana, 35, 840
Cockerell, Theodore D. A., obituary, 166
Codling moth
  aliphatic N-substituted p-Nitrobenzamides against, 151
  control, 347
  baits, processing aromatic chemicals as, 850
  on walnut, 736
Colias philodice eurytheme, 301, 487
Colorado potato beetle, insecticides against, 152
Conotrachelus nenuphar, 7, 12, 19, 514, 632, 774
Corn
  earworm
    control, 933
      studies, 675
    insecticides, 457
  losses and European corn borer, 81
  rootworm
  control, 822
  southern, control of, 908
  seed
    control of *Agonoderus comma* on, 801
    maggot, control of, 908
  wireworm damage reduced by seed treatment, 663
Cotton
  insect control, 551
    in Mexico, 484
  pests, destruction of insect predators during spraying, 904
Cowpea
  curculio control, 856
  insects, control by chlorinated insecticides, 991
Cranberry, new cutworm on, 986
Cruciferae attacked by flea beetles, 144
Cryptococcus fagi, 226
Cucurbits, resistance of, to squash borer, 321
Curculio
  auriger, 825
  proboscideus, 825
Cutworm, new, on cranberry, 986
Cutworms, soil insecticides against, 705
Cyclocephala borealis as a turf pest, 626

DDT
  and *Aëdes* mosquitoes, 835
  analogs, residual toxicity of, to houseflies and mosquitoes, 113
  for *Anopheles* mosquitoes, 447
  for black fly larvae, 392
  for chestnut weevils, 825
  content of milk, 119, 708
  on cows, effect of, on dark rice field mosquito, 393
  deposits and fly strains, 843
  dusts and aphid control on potato, 766
  effect of dispersing and spreading agents upon, 243
  and European fruit *Lecanium*, 147
  failure of, to control house flies, 405
  for flies and roaches, 874
  flood-water mosquitoes, 586
  for fly control, 990
  for horn fly control, 847
  metabolism of, in large milkweed bug, 810
  for *Nomia melanderi*, 519
  oil spray against salt marsh tabanid, 888
  for oriental fruit moth, 25, 351, 786
  for pecan nut casebearer, 357
  permanence of, in powder-post beetle in bamboo, 963
  for potato flea beetles, 470
  recovery of, from treated crops, 602
  residual property of, 420
  residues, new method of evaluating, 151
  resistence to by housefly, 287
  for southern corn rootworm, 992
  spray powders, concentrated, 249
  toxicity of, 611
    to citrus blackfly, 708
    to houseflies, 158
    and sunlight, 154
Dairy cattle, control of mange in, 444
Dalopius pallidus, 942
Dark rice field mosquito, effect on, of feeding on cows treated with DDT, 393
Dendroctonus frontalis, 466
Diabrotica
  duodecimpunctata, 558, 908
  longicornis, 822
Diachus auratus, new strawberry pest, 989
Dicyphus minimus, 983
Diamond back moth in Pa., 681
Dichlorodiphenyl dichloroethane
  for control of flood-water mosquitoes, 586
  recovery of, from treated crops, 602
Dieldrin, wettable powder for fly control, 990
Dinitro compounds and petroleum oil against eggs of European red mite, 44
Dinoderus minutus, 963
Dosage response curve and evaluation time, 579
Drosophila, mass rearing of, 707
Drug store beetle infesting calf starter pellets, 371
Dry-wood termite as a pest, 959
Dusts for corn borer control, 88

Ear tick, life history of, 416
Elm, phloem necrosis of, 729
Enemies, natural, long-tailed mealybug, 777
Entomological Society of Pa., 567
Entomologists
  appraisal of, 1
  army reserve commissions for, 572
  migrations of, 569
Entomology, appraisal of, 1
Entomophagous insects,
  recent introduction of, into California, 995
  technique for evaluating, 546
Eomenacanthus stramineus, 980
Epilachna varivestis, 464, 928, 998
Epitrix hirtipennis, 148
Ethylene dibromide-chlordane mixture against Japanese beetle grubs, 639
European corn borer
  control in Maine, 306
  and dust controls, 88
  and loss in yields, 81
  physiological relationships of, 474
European fruit *Lecanium* in Pa., 853
European red mite eggs, susceptibility to oils and dinitro compounds, 44
Eye-spotted bud moth on prune, 915

Fabric insects, toxicity of insecticides to, 435
Fall army worm control on forage and truck crops, 502
Fig scale control, 48
Fire ant, imported, control of, in Alabama, 976
Flax
  injury by leaf tier, 35
  omnivorous leaf tier on, 840
Flea bettles attacking *Cruciferae*, 144
Fleas, Colorado, recent collections of, 144
Flies
  DDT poisoned, relationship between knockdown and survival time, 874

piperonyl compounds applied to, 423
resistant to DDT, 990
Fluorescent pigments and Japanese beetle studies, 818
Fly
control, 990
repellents for livestock, 281
strains and DDT and methoxychlor deposits, 843
Forage insects and low pressure sprays, 239
Franklinella californica, 920
Fumigant, for seeds, 646
Fumigants
low vapour pressure, relation to toxicity of boiling point, 399
residual action of, 596
Fumigating properties of new insecticides, 436

Gamma-benzine hexachloride
for control of flood-water mosquitoes, 586
toxicity of, 611
Gastroidea cyanea, 613
Gnathocerus cornutus, biology of, 229
Grape
berry moth, control of, 507
mealybug on taxus, 41
Grapes, red-banded leaf roller on, 629
Grapholitha molesta, 25, 786
Grasshopper
control with chlorinated hydrocarbons, 135
populations reduced by gulls, 837
sprays and dusts, comparison of, 956
Green
fruitworm on apricots, 754
peach aphid on tobacco, 677
Gulls reducing grasshopper populations, 837

Harmolita secalis, 154
Heliothis armigera, 457, 675, 904, 933
Hexachlorophenol, acaricidal properties of, 839
Hexaethyl tetraphosphate for cabbage aphid, 156
Honeybee, effect of chemicals on, 855
Honeybees
pollen substitute for, 573
toxicity of
chemicals to, 261, 523
organic insecticides to, 973
Hopkins, A. D., obituary, 868
Horistonotus uhleri, 900
Horn flies
chlorinated hydrocarbon content of milk, 116
control tests and new insecticides, 265
insecticides against, 641
Horn fly
chlorinated hydrocarbons for, 143
control with DDT, 847
Horse flies, biology of, 269
Horse fly
populations, effect of aerial sprays on, 644
repellents, 401
Houseflies
and DDT analogs, 113
effectiveness of chlorinated hydrocarbon residues against, 871
failure of DDT to control, 405
resistant to methoxychlor, 844
resistant strains, 990
temperature and chlorinated hydrocarbon residues, 891
toxicity of
DDT to, 158
parathion to, 990
synthetic compounds to, 532

House fly
effects of weather sanitation, chemical residues, etc., upon, 231
resistance to DDT, 287
Hull, John B., obituary, 869
Hylastinus obscurus, 315
Hylemyia cilicrura, 77
Hypera postica, 554
Hypoderma
bovis, 884
lineatum, 837, 842, 884
Hyppa xylinoides, 986

Imported fire ant control in Alabama, 976
Insect
biting, problems in Alaska, 451
infestations, double sampling of, in evaluation, 396
notes from South America, 761
vectors of cantaloupe mosaic, 770
Insecticidal properties of new chlorinated organic compounds, 127
tests of tropical plants, 549
Insecticide, new chlorinated, for turf-inhabiting insects, 499
Insecticides
against boll weevil, 795
carbon dioxide propelled, toxicity of residues, 123
effect
on peach flavor, 618
of temperature, humidity and temperature on toxicity of, 429
and food law, 564
fumigating properties of, 436
against horn flies, 641
new
effect of on plants and soils, 387
tests to control horn flies, 265
newer, residues of, 966
organic, residual action of, against stable flies, 547
toxicity of, to honeybees, 973
phytotoxicity of vegetables grown in soil treated with, 614
toxicity
to fabric insects and resistance to cleaning, 435
of, to honeybee, 523
Insects
beneficial, transport of,
entomophagous, recently introduced into California, 995
Insemination, artificial, of queen bees, 254

Japanese beetle
baits, phenyl ethyl acetate in, 146
grubs, control of, 639
studies and fluorescent pigments, 818
traps, automatic, 537
Japanese weevil, 540

Kalotermes
hubbardi, 959
minor, 959
Kelly, Edward G., obituary, 162

Lactrodectus mactans, 700
Laphygma frugiperda, 502
Larson, Andrew O., obituary, 165
Larvicides, for red-banded leaf roller on grapes, 629
Lead arsenate, peach twig borer resistant to, 22
Lecanium
corni, 147, 624, 853
nigrofasciatum, 624
Lepidosaphes ficus, 48

Leptinotarsa decemlineata, 152
Leptocoris trivittatus, 994
Limonious agonus, 326
Lindane, 506, 709
  for wireworms, 942
Lipoptena mazamae, record from cattle, 158
Little fire ant, parathion for, 542
Livestock parasite control in Brazil, 276
Long-tailed mealybug, population studies of, 777
Lygus spp. on alfalfa, host plant source of, 93

Macrosiphum
  pisi, 612, 661
  solanifolii, 766
Magicicada septendecim, 359
Mange in dairy cattle, control of, 444
Mealybug
  taxonomy, 835
  wilt and *Pseudococcus brevipes*, 761
Mealybugs, control by *Chrysopa californica*, 560
Melanoplus differentialis, 135, 956
Melanotus communis, 942
Meligethes aeneus in muskmelon breeding, 399
Melitta cucurbitae, 321
Melophagus ovinus, 410, 852, 893, 984
Methoxychlor deposits and fly strains, 843, 845
Methyl bromide
  fumigations of imported orchid plants, 650
  against white-fringed beetle eggs, 733
Mexican
  bean bettle, 996, 998, 999
    control by new insecticides, 928
    studies in California, 464
    toxicity of bis methanes to, 998
  leaf-cutting ant in U. S., 545
  cotton insect control in, 484
Milk
  chlorinated hydrocarbon content of, 116
  DDT content of, 708
Milkweed
  bug, 55
  bug, metabolism of DDT in, 810
  seeds, separation from silks, 55
Mite control on apples, 347
Mites
  control of, 852
    on hothouse tomatoes, 56
Mosquito
  problems in Alaska, 106, 451
  repellents, 686
Mosquitoes
  and DDT analogs, 113
  effectiveness of chlorinated hydrocarbon residues against, 871
  feeding on cows treated with DDT, effect on, 393
  flood-water, larvicides for control of, 586
  radioisotope marking of, 988
  salt-marsh, repellents for, 439
Mounting technique for aphids, etc., 846
Musca domestica, 119, 231, 405, 891
Muskmelon breeding and *Meligethes aeneus*, 399
Myzus persicae, 59, 544, 677, 766

Nabis alternatus, life history of, 991
Nomia melanderi, DDT against, 519

Oil, dormant sprays of, to control peach scales, 624
Olive scale, control by parathion, 656
Omnivorous
  leaf tier, 35
    on flax, 840
Oncopeltus fasciatus, 55, 613, 810
Onion thrips control, 68, 920
Onions, thrips injury to, 756
Orange tortrix
  life history, 37
  host plants, 914
  on red raspberries, control of, 543, 911
Orchid
  insects, 650
  plants, methyl bromide fumigation of, 650
Organic phosphates, insecticidal tests of, 394
Oriental
  beetle as turf pest, 366
  fruit fly in Hawaiian Islands, 713
  fruit moth
    bait jars, 438
    control of, with DDT, 25, 786
    control with DDT and parathion, 351
  insecticides, 348
  parasite, 540
Ornamental plant pests, control of, by parathion, 372
Ornithodoros megnini, 416
Orthosia hibisci, 754
Otobius megnini, 416

Parasite, livestock, control in Brazil, 276
Parasites of long-tailed mealybug, 777
Paratetranchyus pilosus, 624
Parathion
  and bud moth control, 153
  derivatives, toxicity of, 721
  against
    little fire ant, 542
    olive scale, 656
    oriental fruit moth, 351
    pear psylla, 636
    pests of ornamental plants, 372
    pineapple mealybug, 557
  effects on plants, 783
  recovery of, from treated crops, 602
  and red-banded leaf roller, 398
  toxicity
    of, to citrus blackfly, 708
    to flies, 990
Parlatoria oleae, 656
Pea
  aphid, aerosols for control of, 661
  vine silage containing DDT, 119
  weevil, hibernation of, 746
Peach
  control of plum curculio on, 514
  European fruit *Lecanium* on, 147
  flavor, effect of insecticides on, 618
  scales, control by oil in dormant sprays, 624
  insects and disease control, 806
  tree borer control, 343
  twig borer resistant to lead arsenate, 22
Peaches
  benzene hexachloride flavored, 774
  cat-facing on, control of, 335
  plum curculio control on, 330
  plums, injured by red-banded leaf roller, 701
Pear psylla control, 338
  with parathion, 636
Pecan
  minor insect pests of in Texas, 994
  nut casebearer control, 357
Pediculus humanus corporis, 694
Penthaleus major, control of, 852
Periodical cicada, insecticides against, 359
Petroleum oils and dinitro compounds against eggs of European red mite, 44

Phenyl
benzene sulfonates toxicity of, to two-spotted spider mite and Mexican bean beetle, 996
benzoates, toxicity of, to two-spotted spider mite and Mexican bean beetle, 999
ethyl acetate in Japanese beetle baits, 146
Philaenus leucophthalmus, 653
Phloem necrosis of elm, biology of vector, 729
Phlyctaenia rubigalis, 613
Picolinic acid, esters of, as toxicants, 798
Pineapple mealybug, parathion for, 557
Piperonyl
compounds applied to parts of flies, 423
formulations as horse fly repellents, 401
Plum curculio
biology of, 12
control, 330
on peach by chlordan and benzene hexachloride, 514
on prunes, 632
hibernation of, 19
new insecticides against, 7
Plutella maculipennis, 681
Pollen substitute for honeybees, 573
Polychrosis viteana, 507
Popillia japonica, 639, 818
Potato
aphid control on, and DDT dusts, 766
flea beetles, insecticides for, 470
Poultry, absorption of benzene hexachloride in, 980
Powder-post beetle in bamboo, 963
Predators, insect, destruction of by insecticides against cotton pests, 904
Privet, Japanese weevil on, 540
Prune, eye-spotted bud moth on, 915
Prunes, control of plum curculio on, 632
Psallus seriatus, 904
Pseudocneorrhinus bifasciatus, 540
Pseudococcus
brevipes and mealybug wilt, 761
cuspidatae, 41
longispinus, 777
Psorophora confinnis, 393
Psylla pyricola, 338, 636
Pyrausta nubilalis, 81, 88, 474
Pyrethroid, synthetic, effect of commonly used synergists on, 841
Pyrethrum applied to parts of flies, 423

Queen bees artificial insemination of, 254

Radioisotope marking of mosquitoes, 988
Raspberries, red, control of orange tortrix on, 911
control of Willamette mite on, 562
spider mite, tests of insecticides against, 703
Red-banded leaf roller, 29
control, 354
injury to peaches and plums, 701
larvicides against, 629
and parathion, 398
Residual
activity of insecticides against turf insects, 383
property of DDT, 420
Residues, 602
chlorinated hydrocarbon
effect of temperature on, 891
effectiveness against house flies and mosquitoes, 871
on citrus trees, 829
of newer insecticides, 966
toxicity of, 123
Rhagoletis
cingulata, 703
completa, 707
fausta, 703
Rhopalosiphum pseudobrassicae, 73
Rickettsial pox, vector of, 414
Roaches, DDT poisoned, relationship between knockdown and survival time, 874
Rocky Mountain Conference, 567
Rotenone formulations, evaluation of, for cattle grubs, 884
Rye joint worm damage, 154

Salt marsh tabanid control by DDT-oil spray, 888
Sanninoidea exitiosa, 343
Sarcoptes scabiei caprae, 444
Scaphoideus luteolus, 729
Screw-worm survey in Western U. S., 684
Seed-corn
maggot,
biology and control on beans, 77
control of, 908
Seed
stored, insects, silica aerogel against, 553
treatments for wireworms, 942
Seeds
viability of fumigated, 647
Sheep tick
chlorinated hydrocarbons for, 410
control, 852
effect of air temperatures on insecticides against, 893
new insecticides against, 984
pupae, effect of insecticides on, 984
Silica aerogel against stored seed insects, 553
Simulium naevei, 100
Siphona irritans, 116, 265, 641, 847
Sitona cylindricollis, 318
Soil insecticides
against cutworms, 705
for southern corn rootworm, 558
Solenopsis saevissima var. *richteri*, 976
South America, insect notes from, 761
Southern corn rootworm, control of, 558, 992
Southern pine beetle, overlooked relationships of, 466
Soybean flour as pollen substitute for honeybees, 573
Spermatozoon in mealybug taxonomy, 835
Spider mite
two-spotted, 996, 998, 999
and soil fertilization effect on reproduction, 157
Spilonota ocellana, 153, 915
Spittlebug, meadow, insecticides against, 653
Spotted spider mite, toxicity of bis methanes to, 998
Spray conference, Cumberland-Shenandoah, 566
Sprays, low pressure, and forage insects, 239
Spreading agents, effect of DDT upon, 243
Spruce budworm, biology and ecology, 221
Squash borer, resistance of cucurbits to, 321
Stable flies
residual action of insecticides against, 547
use of white mice for testing insecticides against, 461
Stable fly, effects of weather, sanitation, chemical residues, etc., upon, 231
Standard formulations, need for, 403
Stegobium paniceum, 371
Stomoxys calcitrans, 231
Strawberry
crown borer control, 838
new pest of, in Utah, 989
weevil control, 559
Sugar cane, control of wire worms in, 556

Sulfur, interference of, with bioassay dip test, 983
Sweet clover weevil, control and biology, 318
Sweet corn, corn earworm control on, 933
Synergists, commonly used, effect of, on a synthetic pyrethroid, 841
Synthetic compounds, toxicity to house flies of, 532

Tabanid, salt marsh, control by DDT-oil spray, 888
Tabanus
lasiophthalmus, 269
nigrovittatus, 888
phaenops, 271
pumilus, 269
quinquevittatus, 269
sulcifrons, 270
Taxus, grape mealybug on, 41
Termite
dry-wood, as a pest, 959
repellents, 273
Termites, chlordane for, 541
Tetraethyl pyrophosphate dusts, 848
Tetranychus
bimaculatus, 56, 904, 996, 998
willamettei, 562
Texas Entomological Society, 567
Thrips
on asparagus, 849
injury to onions, 756
on tomatoes, 146
onion, control, 68, 920
Thrips tabaci, 920, 68, 756
Ticks
host relationships of, 159
Long Island, host relationships of, 110
Tick repellents, 716, 880
Tobacco
aphids, control, 59, 544, 561
flea beetle control, 148
green peach aphid as pest of, 677
stalk borer in Mexico, 63
Tomatoes, hothouse and mites, 56
Tomato
insects, control of, in California, 666
suckfly, control, 983
thrips infesting, 146
Toxaphene, 225
Toxicants, species-specificity of, 292
Toxicities, comparative, of DDT et al., 611
Toxicity of chemicals to honey bees, 261
Traps, automatic, for Japanese beetles, 537
Trichobaris mucorea, 63
Triethanolamine salt of dinitro-o-sec-butylphenol against fruit tree pests, 380
Tropical plants, insecticidal tests of, 549
Truck crops, fall army worm control on, 502
Turf
inhabiting insects, control of, 499
insects, residual activity of insecticides against, 383
pest
Cyclocephala borealis as, 626
oriental beetle as, 366
Turnip aphids, control, 73

Virus, *polyhedrosis*, against alfalfa caterpillar, 301

Walnut
codling moth on, 736
husk fly, bait trap collection record for, 707
Wasmannia auropunctata, 542
White-fringed beetle eggs, methyl bromide against, 733
White mice, use of for testing repellents and toxicants for stable flies, 461
Willamette mite on red raspberries, 562
Wireworm
control, 327, 942
damage to corn reduced by seed treatments, 663
sand, control in Louisiana, 900
Wireworms, control of, in sugar cane, 556